

The *in vivo* evaluation of antinociceptive and antipyretic activities of *Marrubium deserti* De Noé infusion extract

Saad Somia^{*,†} & Ouafi Saida[†]

Research Laboratory on Arid Zones (LRZA), Faculty of Biological Sciences/University of Sciences and Technology
Houari Boumediene (USTHB), BP n° 32 El-Alia, Bab Ezzouar, 16111 Algiers, Algeria

E-mail: [†]somiasaad89@gmail.com; [†]saida_ouafi@yahoo.fr

Received 05 March 2018; revised 19 March 2019

The aim of this study was to evaluate the *in vivo* antinociceptive and antipyretic activities of *Marrubium deserti* De Noé infusion extract. Antinociceptive effect was evaluated by acetic acid induced writhing response, formalin-induced paw licking and the hot plate method in mice, while antipyretic activity was tested by brewer's yeast induced pyrexia in rats. In each test, we examined the doses of 250, 500 and 1000 mg/kg body weight (bw) of the extract. The infusion extract produced dose-dependent antinociceptive effect against chemically and thermally induced nociceptive pain stimuli in mice. In acetic acid induced writhing test, the inhibition percentage of writhing response was 49.12% at 1000 mg/kg bw ($p < 0.001$). This extract also significantly ($p < 0.001$) inhibited the licking response of the formalin test at 1000 mg/kg bw in both the early phase (64.39%) and the late phase (70.75%). In the hot plate method, the infusion extract significantly ($p < 0.001$) increased the reaction time to heat sensation to 58.21% at 1000 mg/kg bw. Moreover, the extract possessed an excellent ($p < 0.001$) antipyretic effect and even better than the reference drug. These findings indicate that *M. deserti* De Noé infusion extract possesses antinociceptive and antipyretic activities which could be due to the presence of bioactive compounds in this plant.

Keywords: *M. deserti*, Infusion, Phytochemical screening, Acute toxicity, Antinociceptive, Antipyretic

IPC Code: Int. Cl.¹⁹: A01G 13/00, A47G 19/16, A61K 36/00, A62D 101/08, A61K 31/485, A61P 29/00

Pain is defined as an unpleasant sensory and emotional experience associated with tissue damage¹⁻². Analgesics are the substances which decrease pain sensation by increasing pain threshold to external stimuli³. The Non Steroidal Anti-Inflammatory Drugs (NSAIDs) are among the most widely used of all therapeutic agents. They are useful in the treatment of pain and some inflammatory diseases. However, long-term use of the NSAIDs has been associated with gastrointestinal ulceration, bleeding and nephrotoxicity⁴⁻⁵. Therefore, the investigation of new analgesics agents are still a challenge for medical community. As a result more people are turning to herbal medicines for alternative treatment of pain. Herbal drugs have fewer side effects and are largely required as substitute therapeutics⁶.

Traditional medicine is a combination of knowledge and practice, used in diagnosing, preventing and eliminating several diseases. It may rely on past experience and observation handed down from generation to generation verbally or in writing.

Preservation of this information can be a valuable policy for good usage of natural sources. Lamiaceae family has been holding a place of value for hundreds of years due to the numerous aromatic species which used as components of various herbal treatments⁷. The genus *Marrubium* is represented by 97 species which are widely spread over the temperate and warm regions. *Marrubium* species are indigenous in Europe, the Mediterranean area and in Asia⁸⁻¹⁰. Among them, *Marrubium deserti* De Noé known as Merriouet saharai or Djaïdis an endemic herb of central and north Algerian Sahara growing in dry pastures. This plant is a shrub with blancher's leaves and stems. The leaves are velvety and opposite, and are generally terminated by three large teeth of variable forms. The flowers are pale pink¹¹. In Algerian traditional medicine, the leaves and the young buds of *M. deserti* are used in the form of a decoction as a remedy for respiratory diseases like fever, diabetes and as diuretic¹². Furthermore, it is frequently used in traditional medicinal for treating digestive disorders as antispasmodic and as local treatment against scorpion stings and in case of allergy^{10,13}.

*Corresponding author

The biological activities of many *Marrubium* species and their components are reported and it has been showed to possess antinociceptive¹⁴⁻¹⁵, antioxidant¹⁶⁻¹⁹, antimicrobial²⁰, antifungal, cytotoxic²¹, antioedematogenic²², hypolipidemic, hypoglycemic²³, hepatoprotective²⁴, antiproliferative²⁵, vasorelaxant²⁶ and anti-inflammatory effects²⁷.

The essential oil from aerial parts of *M. deserti* De Noé, obtained by hydro-distillation, was analyzed by Gaz Chromatography (GC) and Gaz Chromatography - Mass Spectrometry (GC-MS). Thirty-seven compounds were identified in the oil, with germacrene-D, as the major component (45.7%). This oil was characterized by significant hydrocarbon fraction (78.1%) and the predominance of sesquiterpenes (67.4%)²⁸.

The best known and first isolated diterpenoid was obtained from *M. vulgare* in 1842. It was characterized as marrubiine. This furanien labdane diterpene is, in part, responsible for therapeutic properties of *Marrubium* sp. The genus *Marrubium* is a rich source of diterpenoids, flavonoids and phenylethanoid glycosides²⁹.

The purpose of the present study was to evaluate the antinociceptive and antipyretic activities of the infusion extract from *M. deserti*'s aerial parts in animal models in order to establish a probable mechanism of action and to justify the traditional uses of this plant. The acute toxicity study of the plant was also carried out.

Materials and Methods

Plant Material

Aerial parts of *M. deserti* De Noé were collected from Ain Zaatout town (Biskra, northeast of Algeria) in 2012. The plant species has been identified based on the flora of Quezel and Santa¹¹ and according to the validation of the national institute of the forestry research (INRF), Tamanrasset, Algeria. A voucher specimen of *M. deserti* De Noé (N° 5821-PAM/LRZA/USTHB) has been deposited at the Laboratory of Research on Arid Zones (LRZA), Algiers, Algeria.

The aerial parts were dried and pulverized to powder and preserved for further studies. The infusion extract was prepared according to the traditional method; the fine powder was dissolved in boiling normal saline water (0.9% NaCl). The filtrate was used for biological tests on animals at different doses.

Animals

Wistar rats (*Rattus norvegicus*) (160-200 g) and Albino mice (*Mus musculus*) (25±5 g) of both sexes were procured from Pasteur institute, Algiers, Algeria. Animals were housed in plastic cages at the temperature of 24±2°C and 50±5% relative humidity, with a 12 light/dark cycle respectively. The access to standard pellet diet purchased from the national office of animal feed, Bejaia (ONAB) and water *ad libitum* were allowed for animals. The experiments were carried out in accordance with the current guidelines for the care of laboratory animals, and the ethical guidelines for investigations of experimental pain in conscious animals, as specified by the National Research Council Academies³⁰. The number of animals was the minimum necessary to demonstrate the consistent effects of the drug treatments.

Drugs and Reagents

The chemicals (acetic acid and formalin) used were purchased from Sigma Aldrich GmbH (Steinheim, Germany). Paracetamol and normal saline water (0.9% NaCl) were used as the standard drugs and the control in all studies.

Phytochemical Screening

The phytochemical screening of the infusion extract of *M. deserti* De Noé was performed following the method of Harborne³¹ and Siddiqui *et al.*³². Phytochemical constituents such as flavonoids, terpenoids, saponins, alkaloids, tannins, carotenoids, quinines, lipid and steroids were qualitatively analyzed.

Acute Toxicity of *M. deserti* De Noé in Mice

The acute toxicity study of *M. deserti* De Noé infusion extract was assessed in mice of both sexes, as per OECD-423 guidelines (organization for economic co-operation and development)³³, with slight modifications. Mice were separated into six groups, each consisting of five males and five females mice (n=10). The infusion extract was administered with the doses of 5 g/kg, 7 g/kg, 9 g/kg, 11 g/kg and 13 g/kg bw. The control group received normal saline water at a dose of 20 mL/kg bw. Food and water were withheld 4 h after the drugs administration and the mice were closely observed for the 4 h and then once a day up to 14 days.

Antinociceptive Studies

Acetic Acid Induced Writhing Response in Mice

The analgesic activity was evaluated using acetic acid induced writhing method in mice following the

method of Sawadogo *et al.*³⁴ with slight modification³⁵⁻³⁶. The acetic acid was administered intraperitoneally to animals to create pain sensation. The animals were divided into five groups with six males in each (n=6), they were fasted for 19 h prior to experiments. The groups 1, 2 and 3 were treated with 250, 500 and 1000 mg/kg bw of the infusion extract; while the groups 4 and 5 received normal saline (control) and paracetamol (100 mg/kg bw) as a drug reference respectively. The extract and vehicle were administered orally 30 min before the intraperitoneal injection of the acetic acid (0.6%, v/v in normal saline, 10 mL/kg bw), and the animals were placed individually in transparent plastic cages for counting the number of writhing they made in 10 min beginning just 5 min after their administration. Contraction of the abdominal muscles accompanied by stretching the hind limbs was accepted as writhing movements³⁷. A significant reduction in the number of writhing compared to the control animals was considered as an antinociceptive response³⁸. The analgesic effect was calculated as the inhibition percentage of writhing.

$$\% \text{ inhibition} = (1 - Wt/Wc) \times 100$$

Where Wc is the average writhing response of the control group and Wt is the average writhing response of the treated group.

Formalin Induced Paw Licking in Mice

In this experiment, we used the method of Ullah *et al.*³⁸ with slight modifications³⁹, where pain was induced by formalin. The five groups of mice (n=6); vehicle control (normal saline water, 0.5 mL), paracetamol (100 mg/kg bw) and the infusion extracts of the plant (250, 500 and 1000 mg/kg bw), were administered orally thirty minutes after the subcutaneously injection of 50 µL of a freshly prepared 0.6% formalin solution that was injected under the plantar surface of the right hind paw of each mouse. The time in second spent in licking and biting responses of the injected paw was taken as an indicator of pain response. Antinociceptive effect was determined in two phases. The early phase was recorded during the first 5 min, while the late phase was recorded during the last 15 to 30 min. The inhibition percentage of the paw licks for both phases calculated by following formula:

$$\% \text{ inhibition} = \frac{Do-Dt}{Do} \times 100$$

Where Do is the average number of paw licks of the control group and Dt is the number of paw licks of the treated group.

Hot Plate Test in Mice

The 'hot-plate' test (thermal test) used in this study was described by Lanthers *et al.*⁴⁰ and Ojewole⁴¹ to evaluate the analgesic activity. The temperature of the hot plate was regulated at 55±0.5°C. The five groups of 6 mice each were fasted for 18 h and water was provided *ad libitum*. The mice of each group were placed on the hot plate in order to obtain its response to electrical heat induced pain stimulus. Licking of the paws or jumping out of the hot plate was taken as an indicator of pain response. Reaction time in seconds was taken as the interval between the instant that the animal reaches the hot plate till the moment that the animal licks its paws or jumps out. Each mouse was treated orally with control (normal saline water, 0.5 mL), paracetamol (100 mg/kg bw) or with the infusion extract at doses of 250, 500, 1000 mg/kg bw. Thirty minutes after treatment, the reaction times of each group were evaluated five times individually each hour. The percentage thermal pain stimulus protection was calculated as the formula:

$$\% \text{ protection} = \frac{T_a - T_b}{T_a} \times 100$$

Where Ta is the reaction time (s) after drug administration; Tb is the reaction time (s) before drug administration.

Antipyretic Activity in Rats

Antipyretic activity of *M. deserti* De Noé infusion extract was measured following the method described by Fadeyi *et al.*⁴². Pyrexia was induced in rats by subcutaneously injecting brewer's yeast suspension (20% w/v in normal saline, 10 mL/kg bw) into the rat's dorsum region. The initial rectal temperature of each rat was recorded 19 h before the yeast injection using a thermometer (Hartmann, Germany). The animals were then fasted for all the duration of the experiment, water was available *ad libitum*. Nineteen hours after the injection, the rectal temperature of each rat was recorded again to determine the pyretic response to yeast. Only rats that showed an increase in temperature of at least 0.5°C were selected and then rats were divided into five groups with six each (n=6). The infusion extract at doses of 250, 500 and 1000 mg/kg bw of *M. deserti* De Noé were administered orally, while the control group was given normal saline water and the fifth group was treated with paracetamol (100 mg/kg bw). The temperature was measured at 1, 2, 3 and 4 h after drug and vehicle administrations.

Statistical Analysis

Results are expressed as mean \pm standard deviation (S.D.). Statistical significance was determined using one way ANOVA followed by Tukey's test for multiple comparisons. p -values < 0.05 were considered as significant.

Results

Phytochemical Screening

Phytochemical screening of the infusion extract of *M. deserti* De Noé revealed the presence of various bioactive components of which condensed tannins, catechic tannins, alkaloid, steroids, saponins and terpenoid (Table 1). In addition, we demonstrated the presence of high content of total phenolic compounds such as flavonoid, coumarine and glucosides.

Acute Toxicity

Oral administration of *M. deserti*'s infusion extract (up to 13 g/kg bw) did not show any signs of toxicity or behavioral changes in mice during the initial 4 hours after *M. deserti* De Noé infusion extracts

Table 1 — Phytochemical Analysis of the Infusion Extract of *M. deserti* De Noé Aerial Parts

Extract	Phytochemical constituents	Results
<i>M. deserti</i> 's infusion extract	Flavonoids	++
	Lipoids	-
	Condensed tannins	+++
	Catechic tannins	++
	Alkaloids	+++
	Steroids	+
	Saponins	+++
	Quinines	-
	Terpenoids	+
	Coumarine	+++
	Glucosides	+++

Symbols

'+++' indicates presence in high concentration;
'++' indicates presence in moderate concentration;
'+' indicates presence in low concentration and
'-' indicates absence of phytochemicals.

Table 2 — Effect of the Infusion Extract of *M. deserti* De Noé on Acetic Acid-induced Writhing in Mice

Groups	Dose	Number of writhing	% Inhibition
Control		57 \pm 1.7	-
Paracetamol	100 mg/kg	29 \pm 1.4***	49.12
Infusion extract of	250 mg/kg	41 \pm 2.2*** ξ	28.07
<i>M. deserti</i> De Noé	500 mg/kg	33 \pm 2.09*** ξ	42.10
	1000 mg/kg	29 \pm 1.7*** ξ	49.12

Data represent mean \pm SD, (n=6). (-): no activity, *** $p < 0.001$ significant from the control, $\xi p < 0.05$ significant from the reference drug (Paracetamol) (one way ANOVA followed by Tukey's test).

administration. During the 14 days of experiment, no deaths were observed in any of groups.

Acetic Acid Induced Writhing Response in Mice

The results of the analgesic effects of *M. deserti* De Noé infusion extracts on acetic acid induced writhing test in mice are presented in Table 2. The oral administration of the infusion extract significantly ($p < 0.001$) decreased in a dose-dependent manner the number of writhing induced by acetic acid. The infusion extract at doses of 250 and 500 mg/kg b.w. showed 28.07% and 42.10% inhibition of writhing respectively compared to the control group, whereas at higher dose (1000 mg/kg b.w.) significantly ($p < 0.001$) was found to be the same as the standard drug (Paracetamol, 100 mg/kg bw) with 49.12% inhibition.

Formalin Induced Paw Licking

The effect of *M. deserti*'s infusion extract on formalin induced paw licking in mice is shown in Fig. 1 and Fig. 2. There was a dose-dependent




Fig. 1 — Analgesic activity of the infusion extract of *M. deserti* De Noé aerial parts using formalin induced paw licking test on the early phase Data represent mean \pm SD, (n=6), ** $p < 0.01$ significant compared to control, $\xi p < 0.05$ significant compared to reference (one way ANOVA followed by Tukey's test)




Fig. 2 — Analgesic activity of the infusion extract of *M. deserti* aerial parts using formalin induced paw licking method on late phase Data represent mean \pm SD, (n=6), ** $p < 0.01$ significant compared to control, $\xi p < 0.05$ significant compared to reference (one way ANOVA followed by Tukey's test)

Table 3 — Analgesic Activity of the Infusion Extract of *M. deserti* De Noé Using the Hot Plate Method

Groups	Dose	Reaction time at different time intervals (in s)					
		0 h	30 min	1h	2 h	3 h	4 h
Control		9.263±1.03	7.555±0.5	4.856±0.8	5.431±1.03	3.533±0.5	3.933±0.5
Paracetamol	100 mg/kg	7.843±0.9*	16.028±1.7**	17.635±1.2**	20.445±1.3*	19.958±0.7**	20.063±0.6**
Infusion extract of <i>M. deserti</i>	250 mg/kg	8.425±0.3*	14.513±0.7**	11.44±1.2** [£]	10.616±1.1** [£]	9.12±0.8** [£]	8.706±0.6** [£]
	500 mg/kg	9.168±0.9*	14.601±1.04**	12.62±0.8** [£]	14.686±0.8** [£]	13.16±0.7** [£]	10.293±0.3** [£]
	1000 mg/kg	7.965±1.2*	9.355±0.7** [£]	18.216±0.6**	16.853±0.5** [£]	19.063±1.6**	18.261±1.07**

Values represent mean±SD, (n=6), * $p<0.05$ ** $p<0.01$ significant compared to control, [£] $p<0.05$ significant compared to paracetamol (one way ANOVA followed by Tukey's test).

Table 4 — Effect of *M. deserti*'s Infusion Extract Aerial Parts on Yeast Induced Pyrexia in Rats

Groups	Dose	Before yeast		Time after treatment (h)			
		0 h	19 h	1 h	2 h	3 h	4 h
Control		36.93±0.1	38.08±0.1	38.16±0.3	38.23±0.3	38.46±0.2	38.61±0.3
Paracetamol	100 mg/kg	37.48±0.3	38.46±0.3	37.58±0.3*	37.1±0.4**	37.31±0.3**	36.98±0.3**
Infusion extract of <i>M. deserti</i>	250 mg/kg	37.1±0.1	38.66±0.2	37.55±0.3*	37.33±0.5**	36.93±0.6**	36.68±0.5**
	500 mg/kg	36.6±0.3	38.2±0.2	37.93±0.3*	37.55±0.4**	36.96±0.5**	36.85±0.3**
	1000 mg/kg	36.68±0.3	38.33±0.2	38.05±0.4*	38.1±0.4** [£]	37.46±0.3**	37.48±0.4**

Values represent mean±SD, (n=6). * $p<0.05$, ** $p<0.001$ significant compared to control, [£] $p<0.05$ significant compared to paracetamol (one way ANOVA followed by Tukey's test).

inhibition of both phases in the extract tested of the formalin test. The infusion extract at the doses of 500 and 1000 mg/kg bw significantly ($p<0.001$) inhibited the liking response in both the early phase (46.91%, and 64.39% respectively), and the late phase (57.69% and 70.75% respectively) of the formalin test, while at the dose of 250 mg/kg bw, the infusion extract exerted its action only on the early phase (44.75%) compared with the late phase (17.98%). The standard drug paracetamol (100 mg/kg bw) showed a significant ($p<0.001$) inhibition in both phases, with 59.15% inhibition in the early phase and 76.09% inhibition in late phase.

Hot Plate Test

The results of the effect of *M. deserti* De Noé infusion extract on hot plate induced pain in mice are presented in Table 3 and Fig. 3. The infusion extract produced a dose-related significant ($p<0.01$) analgesic effect that continued for 4 h (58.21%) at the dose of 1000 mg/kg b.w. and until 3 h (37.57%) at the dose of 500 mg/kg bw, but at the low dose of 250 mg/kg bw this extract presented a short analgesic activity (41.94%) against thermal pain stimulus that continued only 30 min after the administration. The paracetamol at the dose of 100 mg/kg bw increased the reaction time of heat sensation in mice with 61.63% protection in the 3rd hour which was significantly greater ($p<0.01$) than that seen with the control group.




Fig. 3 — Protection percentage against thermal pain stimulus of *M. deserti* De Noé infusion extract at different time intervals on hot plate test [£] $p<0.05$ significant compared to paracetamol (one way ANOVA followed by Tukey's test)

Antipyretic Activity in Rats

The effect of *M. deserti* De Noé infusion extracts on yeast induced pyrexia is shown in Table 4. Subcutaneous injection of yeast suspension clearly increased rectal temperature of rats 19 h after yeast injection. The oral administration of the infusion extract of *M. deserti* De Noé at the dose of 250 mg/kg and 500 mg/kg bw after 3 h of treatment caused a significant ($p<0.01$) decrease in temperature compared to control group. In contrast, the higher

dose (1000 mg/kg bw) did not elicit a greater reduction from yeast induced pyrexia as would have been expected. However, the infusion extract at the dose of 250 mg/kg bw showed the highest activity even more than the reference drug (Paracetamol, 100 mg/kg bw).

Discussion

In this study, antinociceptive and antipyretic activities of the infusion extract of *M. deserti* De Noé aerial parts were evaluated in addition to the determination of its acute toxicity profile. The antinociceptive potential was assessed using the chemicals (acetic acid and formalin) and the thermal stimuli models (hot plate test) in mice, while the antipyretic test was evaluated on yeast induced pyrexia in rats. The obtained results showed that the infusion extract of this plant demonstrate high significant ($p < 0.001$) effects in all the models used while the acute toxicity results indicate that the *M. deserti*'s infusion extract is harmless for the medicinal use as the relatively high LD₅₀ value (>13 g/kg) from a single exposure, probably suggests that this plant is safe.

In an attempt to determine whether the infusion extract attenuated either the peripheral or central, or both levels of nociception, acetic acid induced writhing, formalin induced paw licking and thermal induced nociceptive tests were performed.

Acetic acid, the peripheral analgesic agent, causes inflammatory pain by inducing abdominal constrictions and stretching of hind limbs⁴³. Recently, it has been found that this response is mediated by the prostaglandin pathways⁴⁴. Therefore, the agent reducing the number of writhing will provide analgesic effect preferably by inhibition of prostaglandin synthesis, a peripheral mechanism of pain inhibition⁴⁵. Our results showed that the infusion extract of *M. deserti* De Noé in dose-dependent manner caused high significant ($p < 0.001$) reduction in the number of writhes. The ability of the infusion extract of *M. deserti* De Noé aerial parts to attenuate the acetic acid induced writhing response test indicates the presence of analgesic compounds that might influence the prostaglandin pathways.

The formalin test is a model of continuing pain including peripheral inflammation and central sensitization. The injection of formalin into the mouse hind paw initiates triphasic spontaneous nociceptive behaviors consisting of flinching, licking and/or biting

of the injected paw². This method shows a biphasic response, the early phase, classified as a neurogenic pain, is an acute response observed immediately after the administration of formalin and lasts for 5 min (0–5 min) as a result of an immediate action of injected formalin on nociceptors. The late phase, classified as an inflammatory pain, is a tonic response resulting from the inflammatory processes generated by the release of inflammatory mediators such as histamine, serotonin, prostaglandin and bradykinin⁴⁶ and the activation of the neurons in the dorsal horns of the spinal cord which appears between 15 and 60 min after the formalin administration⁴⁷⁻⁴⁸. The results of the present study showed that the infusion extract of *M. deserti* De Noé produced antinociceptive activity against both neurogenic and inflammatory phases of formalin induction which suggests that it contain active analgesic principles acting centrally and peripherally, where the last action is supported by the results recorded in the acetic acid induced writhing test.

Another model of nociception that has been widely used to further support the antinociceptive effect observed in any new compound or extract is the thermal induced nociceptive model or hot plate test. The hot plate test measures the response to a brief, noxious stimulus having a close resemblance to clinical pain. This test measures the complex responses to a non-inflammatory, acute nociceptive impulse and is one of the selective models for studying only the central antinociceptive activity⁴⁹. Drugs that act centrally activate the release of endogenous peptides, which is then carried to the spinal cord to inhibit the pain impulse transmission within the dorsal horn⁵⁰. Based on the ability of the infusion extract to prolong the latency on hot plate, we suggest that this extract possessed centrally mediated antinociceptive activity against the thermal induced nociception.

To investigate the antipyretic effect of *M. deserti*'s infusion extract, the yeast induced pyrexia test in rat was carried out. Antipyretic activity is a characteristic of drugs or compounds which have an inhibitory effect on prostaglandin biosynthesis⁵¹. It is well known that pyretic activity involves stimulation of the region in the hypothalamus that controls body temperature; via prostaglandins synthesized within the central nervous system and that the blood-brain barrier prevents drug molecules or other chemicals from entering the central nervous system⁴⁴. Regulation of body temperature

requires a delicate balance between production and loss of heat, and the hypothalamus regulates the set point at which body temperature is maintained⁵². The infusion extract of *M. deserti* De Noé aerial parts possesses a significant antipyretic effect in yeast-induced temperature in rats. So, we suggest that this extract may exhibit central action because pyrexia is central processes⁴².

Furthermore, the chemical screening for compounds known for their potential activities was investigated. Phytochemical screening of the infusion extract of *M. deserti* De Noé aerial parts indicates the presence of tannins, alkaloids, steroids, terpenoids and a high amount of phenolic compounds. Flavonoids constitute a wide array of biologically active compounds that are found abundantly in plant kingdom and dietary intake. They are reported to be effective against different diseases such as antioxidant, analgesic, anti-inflammatory⁵³⁻⁵⁵, antimicrobial⁵⁶, hepatoprotective⁵⁷ and anticancer²⁵ activities. Furthermore, condensed tannins have been suggested to possess free radical scavenging and antioxidant, anti-inflammatory, and hepatoprotective activities⁵⁸. Besides, alkaloids are well known for their abilities to inhibit pain perception¹. Based on these reports, it is possible that the antinociceptive and antipyretic activities of the infusion extract of *M. deserti* De Noé may be due to the presence and the action of flavonoids, tannins, alkaloids or terpenoids.

Conclusion

To the best of our knowledge, this work reveals, for the first time, that *Marrubium deserti* De Noé infusion extract possesses antinociceptive and antipyretic effects, suggesting that the mechanism of action seems to be similar to NSAIDs drugs such as paracetamol. It has been proved that *M. deserti* De Noé has obvious beneficial effects against centrally and peripherally pain models. The relatively high LD₅₀ value (>13 g/kg bw) obtained for the plant shows that it is safe. These findings justify the use of this plant in the Algerian traditional medicine in the treatment of various types of inflammations and pains. However, the chemical constituents responsible for the pharmacological activities remain to be investigated.

Acknowledgement

This study was supported by General Direction of Scientific Research and Technological Development (DGRSDT) with the Ministry of Higher Education and Scientific Research (MHESR).

References

- 1 Drabu S, Chaturvedi S & Sharma M, Analgesic activity of methanolic extract from the fruits of *Trichosanthes dioica*, *Inter J Res Pharmac Biomed Sci*, 3(1) (2012) 399-405.
- 2 Prabhu VV, Nalini G, Chidambaranathan N & Sudarshan Kisan S, Evaluation of anti-inflammatory and analgesic activity of *Tridax procumbens* Linn against formalin, acetic acid and cfa induced pain models, *Inter J Pharm Pharmac Sci*, 3(2) (2011) 126-130.
- 3 Akriti P & Pawar RS, A study on *Ajuga bracteosa* wall ex. Benth for analgesic activity, *Inter J Curr Biol Med Sci*, 1 (2011) 12-14.
- 4 Mishra D, Ghosh G, Sudhir Kumar P & Kumar Panda P, An experimental study of analgesic activity of selective COX-2 inhibitor with conventional NSAIDs, *Asian J Pharmac Clin Res*, 4(1) (2011) 78-81.
- 5 Zulfiker AHM, Mahbubur Rahman M, Kamal Hossain M, Hamid K, Mazumder MEH *et al*, *In vivo* analgesic activity of ethanolic extracts of two medicinal plants *Scoparia dulcis* L. and *Ficus racemosa* Linn., *Biol Med*, 2(2) (2010) 42-48.
- 6 Meher BR, Jena J & Rath BG, Evaluation of analgesic activity of ethanolic extract of *Sphaeranthus indicus*, *Der Pharmac Lettr*, 3(3) (2011) 357-360.
- 7 Naghibi F, Mosaddegh M, Mohammadi Motamed S & Ghorbani A, Labiatae Family in folk Medicine in Iran: from Ethnobotany to Pharmacology, *Iran J Pharm Res*, 2 (2005) 63-79.
- 8 Meyre-Silva C & Cechinel-Filho V, A review of the chemical and pharmacological aspects of the genus *Marrubium*, *Curr Pharmac Design*, 16(31) (2010) 3503-3518.
- 9 Popoola OK, Elbagory AM, Ameer F & Hussein AA, Marrubiin, *Molecules*, 18 (2013) 9049-9060.
- 10 Neffati N, Aloui Z, Karoui H, Guizani I, Boussaid M *et al*, Phytochemical composition and antioxidant activity of medicinal plants collected from the Tunisian flora, *Nat Prod Res*, 31(13) (2017) 1583-1588.
- 11 Quezel P & Santa S, Nouvelle flore de l'Algérie et des régions désertiques méridionales, Tome II, Centre National de la Recherche Scientifique, France, 1963, 1170.
- 12 Benhammou N, Bekkara FA & Coustard J, Antioxidant activity of methanolic and water extracts from *Marrubium deserti* (de Noé) and *Thymelaea microphylla* from Algerian Sahara, *Adv Food Sci*, 31(4) (2009) 194-201.
- 13 Ould El Hadj MD, Hadj-Mahammed M & Zabeirou H, Place des plantes spontanées dans la médecine traditionnelle de la région d'Ouargla (Sahara septentrional Est), *Courr Sav*, 3 (2003) 47-51.
- 14 De Jesus RA, Cechinel-Filho V, Oliveira AE & Schlemper V, Analysis of the antinociceptive properties of marrubiin isolated from *Marrubium vulgare*, *Phytomedicine*, 7(2) (2000) 111-5.
- 15 Meyre-Silva C, Yunes RA, Schlemper V, Campos-Buzzi F & Cechinel-Filho V, Analgesic potential of marrubiin derivatives, a bioactive diterpene present in *Marrubium vulgare* (Lamiaceae), *Farmaco*, 60(4) (2005) 321-6.
- 16 Laouer H, Yabrir B, Djeridane A, Yousfi M, Beldovini N *et al*, Composition, antioxidant and antimicrobial activities of the essential oil of *Marrubium deserti*, *Nat Prod Commun*, 4(8) (2009) 1133-1138.
- 17 Hamedeyazdan S, Sharifi S, Nazemiyeh H & Fathiazad F, Evaluating antiproliferative and antioxidant activity of *Marrubium crassidens*, *Adv Pharm Bull*, 4(1) (2014) 459-464.

- 18 Stanković MS, Total phenolic content, flavonoid concentration and antioxidant activity of *Marrubium peregrinum* L. extracts, *Kragujevac J Sci*, 33 (2011) 63-72.
- 19 Kadri A, Zarai Z, Békir A, Gharsallah N, Damak M *et al*, Chemical composition and antioxidant activity of *Marrubium vulgare* L. essential oil from Tunisia, *Afr J Biotech*, 10(19) (2011) 3908-3914.
- 20 Bokaeian M, Saboori E, Saeidi S, Niazi AA, Amini-Borojeni N *et al*, Phytochemical analysis, antibacterial activity of *Marrubium vulgare* L against *Staphylococcus aureus* in vitro, *Zahedan J Res Med Sci*, 16(10) (2014) 60-64.
- 21 Zarai Z, Kadri A, Ben Chobba I, Ben Mansour R, Bekir A *et al*, The *in-vitro* evaluation of antibacterial, antifungal and cytotoxic properties of *Marrubium vulgare* L. essential oil grown in Tunisia. *Lipids Health Dis*, 10 (2011) 161.
- 22 Stulzer HK, Tagliari MP, Zampirolo JA, Cechinel-Filho V & Schlemper V, Antioedematogenic effect of marrubiin obtained from *Marrubium vulgare*, *J Ethnopharmacol*, 108(3) (2006) 379-84.
- 23 Herrera-Arellano A, Aguilar-Santamaría L, García-Hernández B, Nicasio-Torres P & Tortoriello J, Clinical trial of *Cecropia obtusifolia* and *Marrubium vulgare* leaf extracts on blood glucose and serum lipids in type 2 diabetics, *Phytomedicine*, 11(7-8) (2004) 561-567.
- 24 Ibrahim FM, Ibrahim AY & Omer EA, Potential Effect of *Marrubium vulgare* L. extracts on CCL4 model induced hepatotoxicity in albino mice, *World J Pharm Sci*, 2(12) (2014) 1664-1670.
- 25 Hamedeyazdan S, Fathiazad F, Sharifi S & Nazemiyeh H, Antiproliferative Activity of *Marrubium persicum* extract in the MCF-7 Human Breast Cancer Cell Line, *Asian Pacific J Cancer Prev*, 13(11) (2012) 5843-5848.
- 26 El Bardai S, Morel N, Wibo M, Fabre N, Llabres G *et al*, The vasorelaxant activity of marrubenol and marrubiin from *Marrubium vulgare*, *Plant Med*, 69 (2003). 75-77.
- 27 Saad S, Ouafi S & Chabane D, Anti-inflammatory and acute toxicity evaluation of aqueous infusion extract obtained from aerial parts of *Marrubium deserti* De Noé growing in Algeria, *Afr J Tradit Complement Altern Med*, 13 (2016) 71-75.
- 28 Edziri H, Ammar S, Groh P, Mahjoub MA, Mastouri M *et al*, Antimicrobial and cytotoxic activity of *Marrubium alysson* and *Retama* grown in Tunisia, *Pakistan J Bio Sci*, 10 (2007) 1759-1762.
- 29 Dendougui H, Seghir S, Belloum Z, Benayache F, Leon F *et al*, A new labdanediterpene and other constituents from *Marrubium deserti* Noe ex Coss, *Rec Nat Prod*, 5 (2011) 300-304.
- 30 NRCA: National Research Council Academies, Guide for the care and use of laboratory animals, (The National Academies Press, Washington DC), 2001, 220.
- 31 Harborne JB, Phytochemical Methods: A guide to modern techniques of plant analysis, (Chapman and Hall, London), 1973, 302.
- 32 Siddiqui S, Verma A, Rather AA, Jabeen F & Meghvansi MK, Preliminary phytochemicals analysis of some important medicinal and aromatic plants, *Adv Biol Res*, 3(5-6) (2009) 188-195.
- 33 OECD: Organization for Economic Co-operation and Development, Guideline-423, adopted on 17th December 2001.
- 34 Sawadogo WR, Boly R, Lompo M, Some N, Lamien CE *et al*, Anti-inflammatory, analgesic and antipyretic activities of *Dicliptera verticillata*, *Inter J Pharmac*, 2(4) (2006) 435-438.
- 35 Ouédraogo N, Lompo M, Sawadogo RW, Tibiri A, Hay AE *et al*, Etude des activités anti-inflammatoire, analgésique et antipyrétique des décoctés aqueux des feuilles et des racines de *Pterocarpus erinaceus* Poir. (Fabaceae), *Phytothérapie*, (2012) 1-7.
- 36 Das B, Ferdous T, Mahmood QA, Hannan JMA, Bhattacharjee R *et al*, Antinociceptive and anti-inflammatory activity of the bark extract of *Plumeria rubra* on laboratory animals, *Europ J Med Plant*, 3(1) (2013) 114-126.
- 37 Collier H, Dinneen L, Johnson C & Schneider C, The abdominal constriction response and its suppression by analgesic drugs in the mouse, *British J Pharmac*, 32(2) (1968) 295-310.
- 38 Ullah HMA, Zaman S, Juhara F, Akter L, Tareq SM *et al*, Evaluation of antinociceptive, *in-vivo* & *in-vitro* anti-inflammatory activity of ethanolic extract of *Curcuma zedoaria* rhizome, *BMC Complem Altern Med*, 14 (2014) 346.
- 39 Adedapo AA, Aremu OJ & Oyagbemi AA, Anti-Oxidant, anti-Inflammatory and antinociceptive properties of the acetone leaf extract of *Vernonia Amygdalina* in some laboratory animals, *Adv Pharm Bull*, 4(2) (2014) 591-598.
- 40 Lanhers MC, Fleurentin J, Mortier F, Vinche A & Younos C, Anti-inflammatory and analgesic effects of an aqueous extract of *Harpagophytum procumbens*, *Planta Med*, 58 (1992) 117-123.
- 41 Ojewole JAO, Antinociceptive, anti-inflammatory and antidiabetic effects of *Bryophyllum pinnatum* (Crassulaceae) leaf aqueous extract, *J Ethnopharmac*, 99 (2005) 13-19.
- 42 Fadeyi OO, Obafemi CA, Adewunmi CO & Iwalewa EO, Antipyretic, analgesic, anti-inflammatory and cytotoxic effects of four derivatives of salicylic acid and anthranilic acid in mice and rats, *Afr J Biotechnol*, 3(8) (2004) 426-431.
- 43 Hasan SMR, Hossain MM, Akter R, Jamila M, Mazumder MEH *et al*, Analgesic activity of the different fractions of the aerial parts of *Commelina benghalensis* Linn, *Int J Pharmacol*, 6(1) (2010) 63-67.
- 44 Zakaria ZA, Ghani ZD, Nor RN, Gopalan HK, Sulaiman MR *et al*, Antinociceptive, anti-inflammatory, and antipyretic properties of an aqueous extract of *Dicranopteris linearis* leaves in experimental animal models, *J Nat Med*, 62(2) (2008) 179-87.
- 45 Ferdous M, Rouf R, Shilpi JA & Uddin SJ, Antinociceptive activity of the ethanolic extract of *Ficus racemosa* Linn. (Moraceae), *Orient Pharm Exp Med*, 8(1) (2008) 93-96.
- 46 Wheeler-Aceto H & Cowan A, Neurogenic and tissue-mediated components of formalin-induced edema: evidence for supraspinal regulation, *Agents Actions*, 34(1-2) (1991) 264-9.
- 47 Hunskaar S & Hole K, The formalin test in mice: dissociation between inflammatory and non-inflammatory pain, *Pain*, 30(1) (1987) 103-114.
- 48 Sani MHM, Zakaria ZA, Balan T, Teh LK & Salleh MZ, Antinociceptive activity of methanol extract of *Muntingia calabura* leaves and the Mechanisms of action involved, *Evidence-Based Complem Altern Med*, (2012) 1-10.
- 49 Ranjit KS, Akm MR, Mesbahuddin A, Sitesh CB, Achinto S *et al*, Bioactive alkaloid from *Sida cordifolia* Linn. with analgesic and anti-inflammatory activities, *Iranian J Pharmacol Ther*, 5 (2006) 175-178.
- 50 Iboronke GF & Ajiboye KI, Studies on the anti-inflammatory and analgesic properties of *Chenopodium ambrosioides* leaf extract in rats, *Int J Pharmacol*, 3 (2007) 111-115.
- 51 Al-Ghamdi MS, The anti-inflammatory, analgesic and antipyretic activity of *Nigella sativa*, *J Ethnopharmac*, 76 (2001) 45-48.

- 52 Gupta M, Mazumder UK, Sambath Kumar R, Gomathi P, Rajeshwar Y *et al.*, Anti-inflammatory, analgesic and antipyretic effects of methanol extract from *Bauhinia racemosa* stem bark in animal models, *J Ethnopharmac*, 98 (2005) 267-273.
- 53 Kaidama WM & Gacche RN, Anti-inflammatory activity of quercetin in acute and chronic phases of inflammation in Guinea Pigs, *Amer J Phytomed Clin Ther*, 3 (2015) 129-136.
- 54 Kim HP, Son KH, Chang HW & Kang SS, Anti-inflammatory plant flavonoids and cellular action mechanisms, *J Pharmacol Sci*, 96 (2004) 229-245.
- 55 Narayana KR, Sripal Reddy M, Chaluvadi MR & Krishna DR, Bioflavonoids classification, pharmacological, biochemical effects and therapeutic potential, *Ind J Pharmacol*, 33 (2001) 2-16.
- 56 Sandhar HK, Kumar B, Prasher S, Tiwari P, Salhan M *et al.*, A Review of Phytochemistry and Pharmacology of Flavonoids, *Inter Pharmac Scientia*, 1(1) (2011) 25-41.
- 57 Tapas AR, Sakarkar DM & Kakde RB, Flavonoids as nutraceuticals: A review, *Trop J Pharm Resear*, 7 (2008) 1089-1099.
- 58 Benhouda A & Yahia M, Toxicity, analgesic and anti-pyretic activities of methanolic extract from *Hyoscyamus albus*' leaves in *albinos* rats, *Int J Pharm Pharm Sci*, 6(3) (2014) 121-127.